

CERTIFICATE OF ANALYSIS

Cellulase "Onozuka" R-10 from Trichoderma viride	Cat.No.:	16419
E.C. 3.2.1.4	Contr.No.:	131089

Parameter	Method	Specification	Result
Molecular weight		ca. 52 000	
Appearance		beige lyophilisate	corresponds
Activities (U/mg)	Cellulase Hemicellulase Protease (DMC) α-Amylase Pectinase	ca. 1 ca. 1 ca. 0.01 ca. 0.8 ca. 0.4	0.8 0.69 0.018 0.93 0.23
Minimum shelf life		5	09/ 2016
Storage (°C)		-0/6	+2 to +8

Unit definitions

Cellulase

1 unit is the amount of enzymatic activity which catalyzes the liberation of 1 μ mol glucose from sodium carboxymethyl cellulose per minute at 40°C, pH 4.5.

Hemicellulase

1 unit is the amount of enzymatic activity which liberates 1 μ mol of reducing groups from beechwood xylan per hour at 37°C, pH 5.5, calculated as xylose.

Protease

1 DMC-unit is that amount of enzymatic activity which catalyzes the cleavage of 1 µequivalent peptide bond from dimethylcasein per minute at 25°C, pH 7.0, expressed in terms of the appearance of new terminal amino groups.

α-Amylase

1 unit is that amount of enzymatic activity which catalyzes the liberation of 1 µequivalent of reducing groups from soluble starch (Zulkowsky) per minute at 25°C, pH 6.0, calculated as maltose.

Pectinase

1 unit is that amount of enzymatic activity which catalyzes the liberation of 1 μ mol of reducing groups from pectic acid per minute at 25°C, pH 4.5, calculated as D-galacturonic acid.

We do not guarantee that the product can be used for a special application.

This document does not release you from performing the standard control upon receipt of incoming goods.

SERVA Electrophoresis GmbH Printing date: 26.09.2013
Quality Control

Dipl.-Ing. (FH) Bernhard Göckel Daniela Lux-Helmstetter

This report has been computer generated and does not contain a signature.