



Glutamate dehydrogenase from bovine liver	Cat.No. :	22904
E.C. 1.4.1.3 solution	Contr.No.:	170356

Parameter	Method	Specification	Result
Molecular weight		ca. 350 000	
Appearance		colorless solution in 50 % glycerol	corresponds
Activity (U/mg protein)	Glutamate dehydrogenase in the presence of ADP	ca. 100	94
Activity in other units (U/mg protein)	not containing ADP	ca. 40	44.6
Protein content (mg/ml)	Lowry	ca. 20	23.1
Minimum shelf life	NA		12.04.2019
Storage (°C)			+2 to +8

Unit definition

1 Unit is that amount of enzymatic activity which catalyzes the reductive amination of 1 μmole of 2-oxoglutarate per minute at 25°C, pH 7.3 in the presence of ADP.

1 Unit causes the transformation of 1 μ mole 2-oxoglutarate per minute at 25 °C and pH 7.3 under assay conditions not containing ADP.

We do not guarantee that the product can be used for a special application. This document does not release you from performing the standard control upon receipt of incoming goods.

SERVA Electrophoresis GmbH Quality Control

Dipl.-Ing. (FH) Bernhard Göckel

Daniela Lux-Helmstetter

Printing date: 12.04.2017