

SPECIFICATION

SPECIFICATION

Cellulase "Onozuka" R-10 from *Trichoderma viride*
Special quality for ELOS analysis by VDLUFA
E.C. 3.2.1.4

Cat.No.: 16418

Parameter	Method	Specification
Molecular weight		ca. 52 000
Appearance		beige lyophilisate
Activities (U/mg)	Cellulase	1.0
	Hemicellulase	ca. 1
	Protease (DMC)	ca. 0.01
	α -Amylase	ca. 0.8
	Pectinase	ca. 0.4
Storage (°C)		+2 to +8

Unit definitions

Cellulase

1 unit is the amount of enzymatic activity which catalyzes the liberation of 1 μ mol glucose from sodium carboxymethyl cellulose per minute at 40°C, pH 4.5.

Hemicellulase

1 unit is the amount of enzymatic activity which liberates 1 μ mol of reducing groups from beechwood xylan per hour at 37°C, pH 5.5, calculated as xylose.

Protease

1 DMC-unit is that amount of enzymatic activity which catalyzes the cleavage of 1 μ equivalent peptide bond from dimethylcasein per minute at 25°C, pH 7.0, expressed in terms of the appearance of new terminal amino groups.

α -Amylase

1 unit is that amount of enzymatic activity which catalyzes the liberation of 1 μ equivalent of reducing groups from soluble starch (Zulkowsky) per minute at 25°C, pH 6.0, calculated as maltose.

Pectinase

1 unit is that amount of enzymatic activity which catalyzes the liberation of 1 μ mol of reducing groups from pectic acid per minute at 25°C, pH 4.5, calculated as D-galacturonic acid.